

A systematic approach introduced novel targets in rectal cancer by considering miRNA/mRNA interactions in response to radiotherapy

Solmaz Khalighfar^{a,b}, Mohammad Reza Kalhori^c, Taghi Amirani^d, Amirhoushang Poorkhani^d, Vahid Khor^d, Ebrahim Esmati^a, Marzieh Lashkari^a, Ali Najafi^{e,1} and Ali Mohammad Alizadeh^{b,f,1,*}

^aRadiation Oncology Research Center, Tehran University of Medical Sciences, Tehran, Iran

^bBreast Disease Research Center, Tehran University of Medical Sciences, Tehran, Iran

^cRegenerative Medicine Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

^dIschemic Disorders Research Center, Golestan University of Medical Sciences, Gorgan, Iran

^eMolecular Biology Research Center, Systems Biology and Poisonings Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran

^fCancer Research Center, Tehran University of Medical Sciences, Tehran, Iran

Received 6 February 2021

Accepted 13 July 2021

Abstract.

BACKGROUND: The discovery of miRNA/mRNA interactions in several biological samples prompted the researchers to explore new biomarkers in tumors.

OBJECTIVE: We aimed to investigate the interactions of miRNA/mRNA in response to radiotherapy in the plasma samples of rectal cancer patients.

METHODS: Five microarray datasets related to cancerous and non-cancerous individuals were first used to construct networks. The databases of Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) were applied to analyze pathway enrichment. The plasma samples were then collected from 55 patients with recently diagnosed rectal cancer and 10 healthy subjects. For radiotherapy courses, the patients have consecutively received 30 sessions of local radiation for six weeks. At last, the expression of selected genes and miRNAs was experimentally measured before and after radiotherapy by qPCR, and the protein levels of the target genes were measured by ELISA assay. We evaluated the therapeutic responses based on the tumor regression grade of the Dworak classification.

RESULTS: We identified 5 up-regulated and 5 down-regulated miRNAs and 8 up-regulated and 3 down-regulated genes of the databases. There was a significant increase in tumor suppressor miRNAs, including miR-101-3p, miR-145-5p, miR-26a-5p, miR-34a-5p, and a significant decrease in oncomiRs, including miR-221-3p and miR-17-5p, after radiotherapy compared to the pre-treatment. Moreover, the up-regulated miR-17-5p and miR-221-5p and the down-regulated miR-101-3p and miR-145-5p were directly related to rectal cancer through the interaction with the Wnt, RAS, PI3K, and TGF- β signaling pathways. An analysis of receiver operating characteristics showed that miRNAs 221, 17, and 23 were response-related in locally advanced rectal cancer patients.

¹ Ali Najafi and Ali Mohammad Alizadeh contributed equally to this work.

*Corresponding author: Ali Mohammad Alizadeh, Cancer Research Center, Tehran University of Medical Sciences, Tehran, P.O.

1419733141, Iran. Tel./Fax: +98 21 6119 2501; E-mail: aalizadeh@sina.tums.ac.ir

CONCLUSIONS: It seems that monitoring the miRNA/mRNA interactions during radiotherapy can be an appropriate diagnostic tool to track the recovery process and respond to standard therapies.

Keywords: Radiotherapy, rectal cancer, gene, protein, microarray, microRNA

1. Introduction

Many studies have investigated radiation's effects on the cells' biological behavior, like cell death, chromosomal aberration, and mutagenesis. Observations have shown that the sensitivity of cells to radiation is different. The cells with a high proliferation or division are more sensitive to radiation than cells with a low proliferation or division [1]. Studies showed that some cells are not capable of DNA damage repair (DDR) caused by radiation, and the damage can directly or indirectly cause cell death. In this setting, microRNA (miRNA) may play an essential role in regulating DDR-related processes and altering tumors' sensitivity to radiation [2]. Therefore, evaluating miRNA expression in irradiated patients can give us helpful information on how tumors can resist or sensitize the radiation [3]. For example, miR-21 has been known to progress in many diseases that may increase malignancy. The miR-21 level increases in tumor cells after radiation [4]. Likewise, Yan et al. showed that the miR-101 expression is significantly associated with poor clinical outcomes in colorectal cancer patients and can inhibit the expression of ataxia telangiectasia mutated (ATM) and DNA-PK κ genes. Therefore, overexpression of this miRNA in tumor cells may increase their sensitivity to radiation [5].

Moreover, He et al. showed that colorectal cancer (CRC) patients with a low serum miR-101 had a more reduced 5-year overall survival than patients with a high serum miR-101 level. Therefore, it might be a valuable marker for diagnosis and prognosis [6]. MiR-145 is another miRNA that exhibits tumor suppressor activity in several cancers, including colon cancer. The overexpression of miR-145 in SW620 and DLD1-SNAI1 cells could sensitize these cells to radiation therapy [7]. In this setting, miR-145 and miR-101 are associated with biological processes such as proliferation, growth, and apoptosis. Yang et al. also found that overexpression of miR-100 could increase the sensitivity of CCL244 cells to radiation. For the first time, they suggested that miR-100 may play an essential role in regulating colorectal tumor cells [8]. Likewise, several studies have shown that many crucial proteins in colorectal cancer signaling, such as WNT, PI3K, EGFR, P53, TIMP, and epithelial-mesenchymal transition (EMT), can control colorectal cancer via miRNA. Zheng et al. observed that

miR-106b overexpression led to cell radioresistance through direct interaction with PTEN and P21 proteins, increasing tumor cells' survival and proliferation. They detected that miR-106b could activate the PI3K/AKT signaling pathway via PTEN inhibition [9,10].

Generally, the discovery of miRNA/mRNA interactions in several biological samples prompted the researchers to explore new biomarkers in tumors [11]. In this context, analysis of relevant miRNA/mRNA interactions may be a suitable strategy for the early detection, monitoring, and prognosis of different types of cancer, including colorectal cancer [12]. Therefore, it appears that changes in miRNA/target gene expression in response to radiotherapy can provide valuable information on how to use adjuvant therapies to improve radiation-based treatments. We aimed to develop novel predictive miRNA/target gene pathways for rectal cancer using a resourceful systematic approach.

2. Materials and methods

2.1. Microarray data analysis

Expression profiles of miRNAs (GSE125961 and GSE112955) and mRNAs (GSE44172, GSE123390: GPL17586, and GSE81986: GPL570) in rectal cancer patients were extracted from the Gene Expression Omnibus (GEO) database (<https://www.ncbi.nlm.nih.gov/geo/>). The GEO2R tool was employed to identify the differentially expressed genes and miRNAs [13]. The differentially expressed miRNAs were defined according to the following criteria: $\log_{2}FC > 1$ and adj P -value < 0.05 .

2.2. Predicted target genes of candidate miRNAs

The target genes of miRNAs were identified using the online predictive programs, including miRmap (<https://mirmap.ezlab.org/app/>), miRWalk2 (<http://zmf.umm.uni-heidelberg.de/apps/zmf/mirwalk2/>), and Targetscan Release 7.0 (<http://www.targetscan.org>).

2.3. GO term and KEGG pathway analysis by the FunRich dataset

The pathway enrichment analyses of gene ontology

(GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases were executed using the FunRich dataset, the software for the functional classification of genes.

2.4. Experimental design and sampling

Fifty-five patients diagnosed with rectal cancer and ten healthy individuals were recruited from October 2016 and over one year. All the included participants were informed about the study protocol, and the written consent was taken. This trial was registered in the Iranian Registry of Clinical Trial (Clinical trial number: IRCT2016072618745N9).

The eligible cases were recruited after reviewing the medical records of patients who had previously been diagnosed with rectal cancer based on pathologic reports. Inclusion criteria comprised the age between 30–70 years, non-metastatic stage II or III rectal cancer patients, Karofsky Performance Status \geq 70 or Eastern Cooperative Oncology Group = 0–1, and no history of familial rectal cancer. Exclusion criteria were the patients with any immunocompromised states such as AIDS, history of viral hepatitis, and abnormal counts of white blood cells. For radiotherapy courses, the patients have consecutively received 30 sessions of local radiation for six weeks (five times, weekly).

Plasma samples were collected from healthy volunteers and patients with rectal cancer before and after radiotherapy. First, approximately 10 ml blood samples were taken from all participant using Vacutainer disposable blood collection tubes. The blood was then centrifuged at 3000 *g* for 5 min to separate the plasma. Moreover, peripheral blood mononuclear cells (PBMCs) were then isolated by the Ficoll-Hypaque technique. At last, the cells were suspended into 90% Foetal Bovin Serum (FBS) 10% Dimethyl sulfoxide (DMSO), and the plasma and PBMCs were preserved at -80°C .

2.5. ELISA assay

VEGF (ab100663, Sensitivity: 10 pg/ml, Range: 8.23–6000 pg/ml), SMAD4 (ab253211, Sensitivity: 52.31 pg/ml, Range: 125–8000 pg/ml), ZEB2 (LS-F13506, Sensitivity: 0.312, Range: 0.312–20 ng/ml), TGFBR2 (MBS7223133, Detection Range: 1.0–25 ng/ml, Sensitivity: 0.1 ng/ml), STAT3 (ab176655, Sensitivity: 15 $\mu\text{g/ml}$, Range: 15–1500 $\mu\text{g/ml}$), NOTCH1 (ab155437, Sensitivity: 20 pg/ml, Range: 28.67–7000 pg/ml), TET2 (abIN6233837, Detection Range:

0.313–20 ng/ml, Sensitivity $<$ 0.188 ng/ml), MYC (ELH-CMYC-1, Detection Range: 0.62–150 ng/ml, Sensitivity: 0.62 ng/ml), PTEN (ab206979, Sensitivity: 39.9, Range: 125–8000 pg/ml), WEE1 (MBS9318404, Detection Range: 3.12–100 ng/ml, Sensitivity: 1.0 ng/ml), and RB1 (MBS2509425, Detection Range: 3.13–200 ng/ml, Sensitivity: 1.88 ng/ml) were measured using ELISA kits under the manufacturer's instructions.

The selected protein levels were determined by a sandwich ELISA as follows: aliquots of 100 μl /well of their primary antibodies (100 μl) were used to coat 96-well plates and incubated overnight at 4°C . The plates were blocked with PBS containing 1% bovine serum albumin (BSA) for 1 hour at room temperature, followed by washing with washing buffer (PBS) containing 0.1% BSA plus 0.05% Tween 20. The supernatants were diluted at 1:4 or 1:2 with PBS and dispensed into the wells. After 2 hours of incubation at room temperature, the plates were washed, and 100 μl of a 1:10000 dilution of the secondary antibody were added to each well. After 2–4 hours of incubation at room temperature, the plates were thoroughly washed. After washing of the unbound antibody, 100 μl of TMB-peroxidase substrate/chromogen solution were added to each well and incubated at room temperature for 10–20 min. The reaction was stopped with 100 μl of 1 M H_3PO_4 . Absorbance at 450 nm was determined by an automated ELISA reader [14].

2.6. Real-time PCR analysis

The quantitative PCR (qPCR) was used to evaluate the expression of miRNA and genes. First, the total RNA was extracted from the plasma and PBMC samples. Then, the plasma (250 μL) and PBMC (500 μL) samples were added to 750 μL TRIzol (Beijing Tiangen Biotech Co., Ltd.) in 2 ml microtubes. After that, RNA extraction was performed according to the manufacturer's instructions. Then, 10 μl of the total RNA was reverse-transcribed in a 20 μl reaction mix using the miRcute miRNA cDNA First-Strand Synthesis kit (Beijing Tiangen Biotech Co., Ltd.) and cDNA Synthesis Kit Manual (TAKARA BIO INC. Cat. 6 30 v.0708) following the manufacturer's recommendations. Finally, the real-time PCR assay was done using an ABI StepOne plus System (Applied Biosystems; Thermo Fisher Scientific, Inc.) and the miRcute miRNA Fluorescence Quantitative Detection kit (Tiangen Biotech Co., Ltd.). The reactions were performed at 94°C for 2 min, followed by 40 cycles of 94°C for 20 s and 60°C

for 40 s. The SYBR Green method (AccuPower Green Star qPCR Master Mix; Bioneer, Korea) was used for qPCR genes. All the PCR reactions were done in triplicates [15,16]. The U6 and B-actin were used as the internal control for the normalization. The fold changes of candidate miRNAs and mRNAs were calculated by equation (2)^(-ΔΔCT) [17]. The primer sequences have been listed in Table 1.

2.7. Evaluation of response to radiotherapy

All the participants underwent surgical tumor resection after neoadjuvant radiotherapy (30 sessions of local radiation for six weeks). The patients were classified based on TNM 8th edition [18], and the pathological responses were evaluated based on the tumor regression grade (TRG) Dworak classification [19]. According to this classification: the patients in grade 0 (TRG0) had not tumor regression, in TRG1; tumor mass was dominantly observed, in TRG2; the patients showed few tumor cells alongside dominant tumor cells; in TRG3, patients had very few tumor cells in fibrotic tissues, and finally in TRG4, no tumor cells expected to be observed.

2.8. Statistical analysis

The statistical analysis was carried out using GraphPad Prism 7.04 (San Diego, CA) and SPSS 18 (IBM, New York, USA) statistical analysis software. The one-sample K-S test was used to evaluate the normality of the data. The *t*-test and one-way ANOVA were used to analyze the data in two and multiple groups, respectively. Moreover, we made a receiver operating characteristic (ROC) curve to ascertain a cut-off for expression of selected miRNAs and choose the cut-off point. It can provide the best sensitivity and specificity to discriminate between respondent and non-respondent patients to assess the potential practicality of selected miRNAs as a predictive tool. The ROC analysis was performed using SPSS version 20 (SPSS Inc., Chicago, Illinois, USA). The descriptive analysis for quantitative data was performed using mean ± SD. The statistical significance was defined as $P < 0.05$.

3. Results

3.1. Identification of differentially expressed miRNAs (DEMs) and differentially expressed genes (DEGs)

The datasets of miRNAs (GSE125961 and GSE112955) and mRNAs (GSE44172, GSE123390: GPL17586,

and GSE81986: GPL570) have included 102 samples of rectal cancers. According to our analysis, 96 miRNAs were found to have differential expression in the samples. About 75 of 1067 DEGs were identified as novel genes. The top 5 up-regulated miRNAs, including miR-17, miR-20a, miR-221, miR-23a, and miR-200c, and the top 5 down-regulated miRNAs, including miR-34a, miR-141, miR-145, miR-26a, and miR-101, were represented in Table 2. Besides, the target genes of the selected miRNAs were represented in Table 3.

3.2. Enrichment analysis of DEGs

To examine the biological functions of 75 DEGs, GO and KEGG analyses were performed in the DAVID database. The enrichment analysis of miRNAs and their target genes included cell apoptosis, migration, and proliferation pathways (Fig. 1). We demonstrated the top 10 enriched pathways in Fig. 1. All of these genes were selected for the qPCR for further validation.

3.3. Demographic data of the participants

The mean age was 57.3 ± 11.5 and 52.3 ± 12.5 years old in the patients and healthy groups. Unlike the tumor stage, there were no significant differences between the patients and healthy groups concerning the baseline characteristics such as sex, height, weight, body mass index (BMI) ($P > 0.05$) (Table 4).

3.4. Verification of the differential expression of the candidate oncomiRs

Fifty-five patients with localized rectal cancer and ten healthy subjects were examined for evaluating the expression levels of miR-17, miR-200c-3p, miR-23a-3p, miR-20a-5p, and miR-221-3p before and after radiotherapy. Our results showed that the expression level of miR-17-5p ($p < 0.0001$), miR-200c-3p ($p < 0.0001$), miR-23a-3p ($p = 0.0001$), miR-20a-5p ($p < 0.0001$), and miR-221-3p ($p < 0.0001$) before radiotherapy was significantly different compared to the healthy subjects. Except for miR-200c-3p ($p > 0.9999$) and miR-20a-5p ($p = 0.7929$), the expression levels of miR-221-3p ($p < 0.0001$), miR-23a ($p < 0.0001$), and miR-17-5p ($p < 0.0001$) were significantly different compared to the pre-treatment after radiotherapy (Fig. 2). There was no significant difference in the expression levels of miR-221-3p ($p = 0.8694$) and miR-17-5p ($p = 0.2159$) after radiotherapy compared to the healthy subjects. On the other hand, the expression levels of miR-20a-5p ($p <$

Table 1
A list of primers for the RT-PCR

Genes/miRNAs	Forward primer	Reverse primer
PTEN	TGGATTCTGACTTAGACTTGACCT	GGTGGGTATGGTCTTCAAAAGG
VEGFA	AGGGCAGAATCATCACGAAGT	AGGGTCTCGATTGGATGGCA
SMAD4	ACGAACGAGTTGTATCACCTGG	TGCACGATTACTTGGTGGATG
WEE1	GGGCAGAAGATGACCACATGA	GCCAAGGGAAATCTGTAGAAGG
ZEB2	GCGATGGTCATGCAGTCAG	CAGGTGGCAGGTCATTTTCTT
RB1	CTGGACGACTTACTGCCATC	TCCAACCGTGGGAATAATGCT
TGFBR2	GCTTTGCTGAGGTCTATAAGGC	GGTACTCCTGTAGGTTGCCCT
STAT3	ATCACGCCTTCTACAGACTGC	CATCCTGGAGATTCTCTACCACT
NOTCH1	CGCTGACGGAGTACAAGTG	GTAGGAGCCGACCTCGTTG
TET2	ATACCCTGTATGAAGGGAAGCC	CTACCCCGAAGTTACGCTTTTC
MYC	CACACCCACAATTCAGGAAGAG	GACGTGCTACAAGGTGGCA
B-actin	CACCATTGGCAATGAGCGGTTT	AGGTCTTTGCGGATGTCCACGT
miR-17-5P	GCCAGAAGGAGCACTTAGGGCA	TGGTGACAGCTGCCTCGGGA
miR-221-3P	TCCAGGTCTGGGGCATGAACCT	GGGTAGCATTGGTGAGACAGCCA
miR-20a-5P	ACACAGCTGGATGCAAACCTGCAA	AACTCCAGCTTCGGCTGTCTG
miR-200c-3P	GGCTGGGGACCTGAGGCGAT	CGGGGGCCCTCGTCTTACCC
miR-23a-3P	CCAGGCACAGGCTTCGGG	GAACGGAGGGCAACCTA
miR-101-3P	ACAACATGGCTGCACCAACA	TTAATATTTTCAGCTTACCAC
miR-145-5P	ACAAGGTGGGAGCGAGTGGC	CATCCGGCGCGTGTGGCA
miR-141-3P	CCCCATCCAGAGGGGTGAAGG	GGCTCCCGCGTGGTCTCT
miR-26a-5P	GCACATACTAAGGAGCCAAG	TGCCTTTCTCAGCAACTCC
miR-34a-5P	TGAGGGCGGCTGGGAAAGTG	TTCTCCAGCCAAAAGCCGCC
U6	ATGCAGTCGAGTTTCCACAT	CCATGATCACGAAGGTGGTTT

Table 2
The candidate miRNAs in rectal cancer

miRNAs	Adjusted <i>p</i> -value
UP-regulate	
miR-17	1.00E ⁻⁰⁷
miR-20a	4.03E ⁻⁰⁴
miR-200c	1.06E ⁻⁰³
miR-221	6.14E ⁻⁰³
miR-23a	4.33E ⁻⁰¹
Down-regulate	
miR-141	2.34E ⁻²
miR-145	0.01617
miR-26a	0.02410
miR-101	0.01018
miR-34a	0.00264

Table 3
The candidate genes in rectal cancer

Up-regulated	AKT1, EGFR, IGF1R, MET, TGFBR2, BCL2, ACVR1B, MAPK1, MAPK9, MYC, SMAD4, MAP2K1, FZD9, FZD4, FZD6, FZD10, MAP2KR1, KRAS, MAPK, TGFB2, IGF1R, PI3KR1, IGFB1, IGFB2, IGFB2, APPL1, MAPK3, AKT2, STAT3, BRAF, GRB2, TET2, JUN, NOTCH1, DVL1, LEF1, BIRC5, ZEB2, VEGFA
Down-regulated	APC, MSH2, MSH6, TP53, MSH3, TCF7L2, AXIN2, CTNNB1, PTEN, FOXO3, PDCD4, WEE1, E2F2, TCF7, BAK1, APC2, E2F1, GSK3B, RB1, SP1, RB1, GSK3B, SP1, AP1, TCF7, FRAT2, FRAT1, ESR1, ESR2, BAK1, DDB2, E2F3, ESRRG

Table 4
Demographic characteristics of the participants in the present study

Variables	Patients	Control	<i>P</i> -value
Sex (%)			
Male	30 (59.7)	6 (60)	
Female	25 (40.3)	4 (40)	
Age (years) (mean ± SD)	57.3 ± 11.5	52.3 ± 12.5	0.2
Height (Cm) (mean ± SD)	170.9 ± 7.5	168.7 ± 7.4	0.9
Weight (Kg) (mean ± SD)	76.4 ± 7.5	79.4 ± 10.4	0.7
BMI (kg/m ²) (mean ± SD)	27.5 ± 4.0	28.2 ± 4.4	0.8
Tumor stage (%)			
Stage 2	26 (47.3)		
Stage 3	29 (52.7)		0.4

BMI: Body mass index.

0.0001), miR-23a-3p ($p = 0.0002$), and miR-200c-3p ($p < 0.0001$) were significantly different after radiotherapy compared to the healthy subjects, indicating fewer effects of radiotherapy on the expression levels of these miRNAs (Fig. 2).

3.5. Verification of the differential expression of the candidate tumor suppressor miRNAs

The expression levels of miR-26a-5p ($p < 0.0001$), miR-101-3p ($p < 0.0001$), miR-145-5p ($p < 0.0001$), miR-34a-5p ($p < 0.0001$), and miR-141-3p ($p < 0.0001$) were significantly different before radiotherapy

272
273
274
275
276
277
278
279
280
281
282




Fig. 1. Pathways enrichment analysis related to derived miRNAs and their target genes.

283 compared to the healthy subjects. Except for miR-141-
 284 3p ($p > 0.9999$), the expression levels of miR-26a-5p
 285 ($p < 0.0001$), miR-101-3p ($p < 0.0001$), miR-145-
 286 5p ($p < 0.0001$), and miR-34a-5p ($p < 0.0001$) were
 287 significantly different after radiotherapy compared to
 288 the pre-treatment, indicating the positive effects of the
 289 radiotherapy on the expression levels of these miRNAs
 290 (Fig. 3).

291 3.6. Confirmation of selected oncoproteins and tumor 292 suppressor proteins by ELISA

293 The results showed that the expression levels of onco-
 294 proteins, including VEGF, SMAD4, ZEB2, TGFBR2,
 295 STAT3, NOTCH1, TET2, and MYC, were significantly
 296 increased in the cancerous samples compared to the
 297 normal group. Likewise, a similar trend was consistent
 298 as the advanced stage of disease (Table 5). In contrast,




Fig. 2. The expression level of the selected miRNAs, miR-17-5p (a), miR-221-3p (b), miR-20a-5p (c), miR-200C-3p (d), and miR-23a-3p (e), before and after radiotherapy in rectal cancer patients compared to the healthy subjects. The relative expression of selected miRNAs was normalized using U6 as the reference RNA. * $P < 0.05$ is considered as a significant level.




Fig. 3. The selected miRNAs' expression, miR-141-3p (a), miR-145-5p (b), miR-26a-5p (c), miR-101-3p (d), miR-34a-5p (e), before and after radiotherapy in rectal cancer patients compared to the healthy subjects. The relative expression of selected miRNAs was normalized using U6 as the reference RNA. * $P < 0.05$ is considered as a significant level.

Table 5

The protein levels of selected tumor suppressor and oncoproteins in rectal cancer patients in response to radiotherapy

	Normal	Pre-rad	Post-rad
VEGFA (pg/ml)	88.3 ± 9.5	339 ± 47*	98.3 ± 17#
SMAD4 (pg/ml)	134 ± 21	456 ± 49*	143 ± 16#
ZEB2 (ng/ml)	5.3 ± 2.0	17 ± 4.0*	7.0 ± 5.0#
TGFBR2 (ng/ml)	6.5 ± 2.0	21 ± 3.0*	7.6 ± 1.6#
STAT3 (μg/ml)	56 ± 5.0	135 ± 32*	63 ± 19#
NOTCH1 (pg/ml)	115 ± 12	444 ± 39*	125 ± 21#
TET2 (ng/ml)	1.5 ± 0.3	16 ± 3.0*	2.1 ± 1.2#
MYC (ng/ml)	21 ± 6	132 ± 26*	27 ± 11#
PTEN (pg/ml)	888 ± 48	416 ± 33*	834 ± 44#
WEE1 (ng/ml)	76 ± 12	26 ± 10*	68 ± 15#
RB1 (ng/ml)	166 ± 15	75 ± 11*	157 ± 23#

* $P < 0.05$ compared to the normal group. # $P < 0.05$ compared to the pre-treatment group.

the expression levels of tumor suppressor proteins, including PTEN, WEE1, and RB1, were significantly decreased in the advanced stages compared to the primary stages and the normal subjects (Table 5).

3.7. Verification of the differential expression of the candidate genes

We examined the expression levels of the VEGF, SMAD4, ZEB2, TGFBR2, STAT3, NOTCH1, TET2, and MYC oncogenes in the patients before and after radiotherapy compared to the healthy subjects (Fig. 4). Our results showed that the expression levels of these genes were significantly different before radiation from that of the normal subjects. Similarly, there was a significant difference in their expression levels after radiotherapy compared to the pre-treatment (Fig. 4a).

Finally, we have compromised the expression levels of PTEN, WEE1, and RB1 tumor suppressor genes before and after radiotherapy compared to the healthy subjects. The expression levels of these genes were significantly different before radiotherapy from that of the normal subjects. Their expression levels were very different after radiotherapy than the pre-treatment, indicating the positive effects of radiotherapy on these tumor suppressor genes (Fig. 4b).

3.8. Expression evaluation of miRNAs in response to radiotherapy

The ten miRNAs were studied among the participants to validate their potential role as a predictor of response to radiotherapy. The patients were categorized into TRG3/4 and TRG1/2 groups as respondents and non-respondents, respectively. Interestingly, the patients who were in TRG3/4 (65.4%) group had significantly

lower miR-221 ($p = 0.019$), miR-17 ($p = 0.025$), and miR-23 ($p = 0.038$) expression levels. In contrast, they showed significantly higher miR-26 expression levels ($p = 0.005$). No significant differences were observed for the other miRNAs.

ROC analysis was performed to investigate the potential utility of miR-221, miR-17, miR-23 and miR-26 as predictive biomarkers of response to radiotherapy. The AUC value for expression of miR-221, miR-17, and miR-23 was 0.717 (95% CI = 0.552–0.882; $p = 0.009$), 0.695 (95% CI = 0.549–0.842; $p = 0.018$) and 0.659 (95% CI = 0.513–0.806; $p = 0.054$), respectively (Fig. 5a).

Moreover, miR-221 and miR-17 provided a better predictive profile with an AUC value of 0.795 (95% CI: 0.651–0.920) (Fig. 5b). For the miR-26, the AUC was 0.735 (95% CI = 0.595–0.894) (Fig. 5c). In the optimum truncation point, the sensitivity and specificity were 86.1% and 57.9% for miR-221, 69.4%, and 68.4% for miR-17. For the combination of both oncomiRs, the sensitivity and specificity were 66.7% and 84.2%, respectively. The analysis of predictive power of radiotherapy for the miR-26 represented the sensitivity and specificity of 75% and 73.7%, respectively.

4. Discussion

Our results showed the tumor suppressor miRNAs' expression, including miR-101-3p, miR-145-5p, miR-26a-5p, and miR-34a-5p, and also the expression of oncomiRs, including miR-221-3p and miR-17-5p, changed significantly after radiotherapy compared to the pre-treatment in the rectal cancer patients. Moreover, there was a significant difference in the expression level of the oncoproteins and the tumor suppressor proteins after radiotherapy compared to the pre-treatment. The analyses also showed that the up-regulated miR-17-5p and miR-221-5p and the down-regulated miR-101-3p and miR-145-5p were directly related to rectal cancer via the Wnt, RAS, PI3K, and TGF- β signaling pathways.

Previous studies have shown that miRNAs could predict and determine treatment response to cancer treatments. They can also increase the sensitivity of tumors to radiation by inhibiting target genes. In this setting, miRNAs may modify current therapeutic strategies and make them more targeted and effective. Therefore, miRNAs are not only biomarkers involved in cancer prediction, prognosis, diagnosis, and monitoring, but also they can be used as therapeutic targets in many can-




Fig. 4. The expression of the selected oncogenes (a) and tumor suppressor genes (b) before and after radiotherapy in rectal cancer patients compared to the healthy subjects. The relative expression of genes was normalized by using b-actin as the internal control gene. * $P < 0.05$ compared to the healthy group. # $P < 0.05$ compared to the pre-treatment group.




Fig. 5. Receiver operating characteristic (ROC) curve to evaluate the potential efficacy of (a) miR-221, miR-17, and miR-23, (b) the combination of miR-221 and miR-17, and (c) miR-26 to differentiate patients with response to radiotherapy (TRG 3/4) from non-responsive (TRG 1/2). TRG: tumor regression grade.

379 cers [20,21]. Our results showed that changes in miRNA
380 expression in response to tumor radiation could provide
381 helpful information on using adjuvant therapies
382 to improve radiation-based treatments [22]. In this setting,
383 four main pathways can be activated by the growth
384 factor receptors in response to radiation.

385 The first pathway is PI3K/AKT, which modifies the
386 expression of BAX, BIM, BCL2, and FOXO in response
387 to radiation. PTEN is a tumor suppressor protein that
388 plays a crucial role in regulating this pathway. Our
389 results showed that the PTEN level was significantly
390 different after radiotherapy compared to the pre-
391 treatment. In this setting, the miR-21 has been shown
392 to target and inhibit PTEN protein. This miRNA plays
393 a central role in the occurrence or progression of cancer
394 and may increase malignancy [23]. Moreover, Zheng
395 et al. investigated the role of miR-106b in the sensitivity
396 of human colorectal cells to radiotherapy. They observed
397 that the increased expression level of miR-106b

398 led to radiation resistance through direct interaction
399 with PTEN and P21 proteins and improved cell survival
400 and proliferation under radiation. They also observed
401 that miR-106b could activate the PI3K/AKT signaling
402 pathway by restricting PTEN protein and enhancing
403 cell proliferation [9]. Similar to our results, Drebber et
404 al. observed that the expression levels of miR-21 and
405 miR-145 were increased and decreased, respectively,
406 in tumor tissues. Their observations indicated reduced
407 miR-21 and an increase in miR-145 after treatment in
408 rectal tumors [4]. Thus, these miRNAs can be used to
409 monitoring patients with radiotherapy.

410 The second pathway involves MAPK that promotes
411 cell proliferation controlled by RAS and RAF activa-
412 tions. Our results showed that the expression levels of
413 miR-17-5p, miR-200c-3p, miR-23a-3p, miR-20a-5p,
414 and miR-221-3p were significantly different after radio-
415 therapy than the pre-treatment. Moreover, high levels of
416 miR-17-3p expression were related to a shorter disease-

free survival [24]. Likewise, elevated plasma levels of miR-221 could be used as a potential biomarker to predict poor overall survival in CRC patients [25]. Similar to our results, Let-7 [26], miR-145 [27], and miR-143 have been identified as tumor suppressors that can inhibit RAS and RAF expressions [28]. Studies have also shown that the Let-7 family can increase tumor cell sensitivity to radiation by decreasing the RAS family genes [29]. According to our results, the up-regulation of miR-143 and miR-145 may improve the rectal cells' radiosensitivity. Thus, these miRNAs may be used as biomarkers to predict therapy responses.

The third pathway is the transformation of epithelial-mesenchymal transition cells, leading to cancer cell metastasis. Several miRNAs are involved in EMT regulation, such as the miR-200 family. Our results showed that the expression level of miR-200c-3p was significantly different after radiotherapy compared to the pre-treatment. Sun et al. found that miR-429, a member of the miR-200 family, was significantly down-regulated in colon cancer [30]. According to our results, Hur et al. found that miR-200 was abnormally expressed in metastatic colon tumors correlated with reducing the expression of the target genes, such as ZEB1, ETS1, and FLT1 genes. This issue could up-regulate E-cadherin and down-regulate Vimentin sequentially, leading to the EMT signaling pathway [31]. Observations have also shown that factors such as ZEB1 have an active role in the EMT pathway through the interaction of miR-200 and miR-141 [10]. Therefore, the down-regulation of miR-200 might improve the radiosensitivity of the colorectal cells.

The fourth pathway is the P53, identified as a mutated tumor suppressor in 50–75% of all colorectal cancers. P53 can induce several miRNAs' expression and maturation, including Let-7a, miR-133a, miR-34, and miR-16 in colon cancer cells [32]. Our results showed that the expression levels of miR-26a-5p, miR-101-3p, miR-145-5p, miR-34a-5p, and miR-141-3p were significantly different after radiotherapy compared to the pre-treatment. Similar to our results, the miR-34 family can play a central role in the cell cycle, proliferation, apoptosis, and angiogenesis, which targets the CDK4/6, cyclin E2, E2F2, BCL2, and SIRT proteins. Moreover, the miR-34 expression is decreased in colorectal cancers, which may be due to the deletion of 1p36 or miR-34 promoter's methylation [32]. Therefore, the increased expression of miR-34 and Let-7a in tumor cells can increase the sensitivity to radiotherapy.

Overall, our results provided helpful information on inhibiting the expression of proteins involved in

miRNAs' cancer-related pathways. Although some of these interactions are only predicted, this dual computational approach can provide critical information for conducting new validation studies. We showed that over-expressed miRNAs, such as miR-221-3p, and down-expressed miRNAs, such as miR-101-3p and miR-26a-5p, could interact more genes in the signaling pathways. Generally, our observations may hypothesize that miRNAs can enhance proliferation and inhibit cell death. These results can support future studies that may determine the sensitivity, specificity, and efficacy of selected miRNAs, representing particular interactions with genes and molecular pathways in rectal cancer.

Additionally, the previous studies showed that patients with the complete response could undergo less invasive strategies such as “wait and watch” approaches [33]. The supporting literature indicated that such an approach could increase the quality of life in patients who were responsive to neoadjuvant chemoradiotherapy [33]. Contrarily, the patients with resistance to neoadjuvant chemoradiotherapy should manage properly based on their response rate to the treatments. This issue can result in modifying the irradiation dose or administration of different chemotherapeutic agents. Likewise, understanding the underlying mechanisms for resistance can improve the efficacy of radiation therapy by overcoming radioresistance. In the current work, we observed that from 10 selected miRNAs, four associated with the response to radiotherapy. In this respect, we observed that the miR-221, -17, and -23 were response-related miRNAs in patients with locally advanced rectal cancer. This profile has not been previously evaluated as a tool for treatment monitoring in locally advanced rectal cancer to the best of our knowledge. However, some other novel miRNAs were reported by which the response to chemoradiotherapy was predicted [34,35]. In this respect, it was demonstrated that the miR-31 was associated with poor overall survival and a higher rate of resistance in locally advanced rectal cancer [34]. Besides, it was demonstrated that post-surgical expression of miR-345 was more elevated in rectal cancer patients resistant to chemoradiotherapy [36].

Moreover, Campayo et al. reported that the overexpression of miR-21, miR-99-b, and miR-375 was observed among the patients who showed a poor response rate to the chemoradiotherapy. Consistently, they provided similar findings regarding the possible role of let-7b in enhancing radiosensitivity in rectal cancer patients [37]. Interestingly, we observed that post-surgical higher expression of tumor suppressor miRNAs, miR-26, was associated with better radiosensitivity. This is

sue can imply that the target pathways of these miRNAs can be a possible target for targeted therapy for improving the efficacy of chemoradiotherapy.

4.1. Limitation and clinical application

Although we assessed a panel of the biomarkers to predict the response to treatment in locally advanced rectal cancer patients, our low sample size may diminish the validity of the findings. The clinical decision on the management of rectal cancer is substantially made based on the different variables before initiation of any clinical intervention. However, the current decision-making system is not accurate enough. As a result, around 30% of the patients showed no clinical response to performed neoadjuvant interventions [38]. About 70 miRNAs are associated with treatment resistance in rectal cancer patients [39,40]. The possible valuable role of miRNAs in the clinic can be range from the biomarker to candidate for the targeted therapy. So far, multiple techniques developed to facilitate the clinical applications of miRNAs in patient management. These procedures can comprise antisense oligonucleotides or antagomirs, locked nucleic acids, peptide nucleic acids, the newest miRNA sponges and miRNA masking techniques, and an increased tumor suppressor miRNAs by miRNA mimics or viral vector-encoded miRNA replacement [40]. Despite their considerable impact on patient management, these techniques have been studied on a limited scale of rectal cancer patients. We could not find any active clinical trial regarding using these techniques on patients with locally advanced rectal cancer based on our search. Therefore, more studies should be conducted to assess the impact of these strategies in managing radioresistance patients.

5. Conclusion

We indicated that the interactions of selected miRNAs and target genes were associated with cell apoptosis, migration, and proliferation that can play crucial roles in rectal cancer. They may also be new candidate biomarkers to monitor conventional radiotherapy.

Abbreviation

DDR DNA damage repair;
miRNA microRNA;
ATM ataxia telangiectasia mutated;

CRC colorectal cancer;
EMT epithelial-mesenchymal transition;
GEO Gene Expression Omnibus, GO, gene ontology;
KEGG Kyoto Encyclopedia of Genes and Genomes;
PBMCs peripheral blood mononuclear cells;
FBS Foetal Bovin Serum;
DMSO Dimethyl sulfoxide;
BSA bovine serum albumin;
qPCR quantitative PCR;
TRG tumor regression grade;
ROC receiver operating characteristic;
DEMs differentially expressed miRNAs;
DEGs differentially expressed genes;
BMI body mass index.

Acknowledgments

Research reported in this publication was supported by Elite Researcher Grant Committee under award number (Grant Number: 982679) from the National Institute for Medical Research Development (NIMAD), Tehran, Iran, and the Tehran University of Medical Sciences (Grant Number: 32359).

Author contributions

Conception: SK, MRK, VK, and AMA
Interpretation or analysis of data: SK, MRK, ML, AN, and AMA
Preparation of the manuscript: SK, MRK, and AMA
Revision for important intellectual content: TA, AP, and EE
Supervision: AMA

Funding

None of the funding sources had any role in the study design, the collection, analysis, and interpretation of data, or the decision to submit the article for publication.

Compliance with ethical standards

Conflict of interest

The manuscript authors have no conflicts of interest to declare and are responsible for the paper's content.

Ethical approval

All procedures performed in the studies involving animal participants were under the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The study was also conducted under relevant national and international guidelines and approved by the Institutional Animal Care and Use Committee of Tehran University of Medical Sciences.

References

- [1] A.P. Møller and T.A. Mousseau, Biological indicators of ionizing radiation in nature, in: *Environmental Indicators*, Springer, 2015, pp. 871–881.
- [2] F. Cellini, A.G. Morganti, D. Genovesi, N. Silvestris and V. Valentini, Role of microRNA in response to ionizing radiations: Evidences and potential impact on clinical practice for radiotherapy, *Molecules* **19** (2014), 5379–5401.
- [3] M. Agostini, S. Pucciarelli, F. Calore, C. Bedin, M. Enzo and D. Nitti, miRNAs in colon and rectal cancer: A consensus for their true clinical value, *Clinica Chimica Acta* **411** (2010), 1181–1186.
- [4] U. Drebber, M. Lay, I. Wedemeyer, D. VALLböHMER, E. Bollschweiler, J. Brabender, S.P. Mönig, A.H. Hölscher, H.P. Dienes and M. Odenthal, Altered levels of the onco-microRNA 21 and the tumor-suppressor microRNAs 143 and 145 in advanced rectal cancer indicate successful neoadjuvant chemoradiotherapy, *International Journal of Oncology* **39** (2011), 409–415.
- [5] D. Yan, W.L. Ng, X. Zhang, P. Wang, Z. Zhang, Y. Mo, H. Mao, C. Hao, J.J. Olson and W.J. Curran, Targeting DNA-PKcs and ATM with miR-101 sensitizes tumors to radiation, *PLoS One* **5** (2010), e11397.
- [6] D. He, Z. Yue, G. Li, L. Chen, H. Feng and J. Sun, Low serum levels of miR-101 are associated with poor prognosis of colorectal cancer patients after curative resection, *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research* **24** (2018), 7475.
- [7] Y. Zhu, C. Wang, S.A. Becker, K. Hurst, L.M. Nogueira, V.J. Findlay and E.R. Camp, miR-145 antagonizes SNAIL-mediated stemness and radiation resistance in colorectal cancer, *Molecular Therapy* **26** (2018), 744–754.
- [8] X.-D. Yang, X.-H. Xu, S.-Y. Zhang, Y. Wu, C.-G. Xing, G. Ru, H.-T. Xu and J.-P. Cao, Role of miR-100 in the radioresistance of colorectal cancer cells, *American Journal of Cancer Research* **5** (2015), 545.
- [9] L. Zheng, Y. Zhang, Y. Liu, M. Zhou, Y. Lu, L. Yuan, C. Zhang, M. Hong, S. Wang and X. Li, MiR-106b induces cell radioresistance via the PTEN/PI3K/AKT pathways and p21 in colorectal cancer, *Journal of Translational Medicine* **13** (2015), 252.
- [10] O. Slaby, M. Svoboda, J. Michalek and R. Vyzula, MicroRNAs in colorectal cancer: Translation of molecular biology into clinical application, *Molecular Cancer* **8** (2009), 102.
- [11] A. Baldassarre, C. Felli, G. Prantera and A. Masotti, Circulating microRNAs and bioinformatics tools to discover novel diagnostic biomarkers of pediatric diseases, *Genes* **8** (2017), 234.
- [12] A. Polo, A. Crispo, P. Cerino, L. Falzone, S. Candido, A. Giudice, G. De Petro, G. Ciliberto, M. Montella and A. Budillon, Environment and bladder cancer: Molecular analysis by interaction networks, *Oncotarget* **8** (2017), 65240.
- [13] J. Hu, G. Cai, Y. Xu and S. Cai, The plasma microRNA miR-1914* and-1915 suppresses chemoresistant in colorectal cancer patients by down-regulating NFIX, *Current Molecular Medicine* **16** (2016), 70–82.
- [14] V. Khorii, A.M. Alizadeh, S. Khalighfard, Y. Heidarian and H. Khodayari, Oxytocin effects on the inhibition of the NF- κ B/miR195 pathway in mice breast cancer, *Peptides* **107** (2018), 54–60.
- [15] S. Khalighfard, A.M. Alizadeh, S. Irani and R. Omranipour, Plasma miR-21, miR-155, miR-10b, and Let-7a as the potential biomarkers for the monitoring of breast cancer patients, *Scientific Reports* **8** (2018), 17981.
- [16] S. Farsinejad, M. Rahaie, A.M. Alizadeh, M. Mir-Derikvand, Z. Gheisary, H. Nosrati and S. Khalighfard, Expression of the circulating and the tissue microRNAs after surgery, chemotherapy, and radiotherapy in mice mammary tumor, *Tumor Biology* **37** (2016), 14225–14234.
- [17] V. Khorii, A.M. Alizadeh, Z. Gheisary, S. Farsinejad, F. Najafi, S. Khalighfard, F. Ghafari, M. Hadji and H. Khodayari, The effects of low-level laser irradiation on breast tumor in mice and the expression of Let-7a, miR-155, miR-21, miR125, and miR-376b, *Lasers in Medical Science* **31** (2016), 1775–1782.
- [18] M.F. Amin, F.L. Greene, S.B. Edge, C.C. Compton, J.E. Gershenwald, R.K. Brookland, L. Meyer, D.M. Gress, D.R. Byrd and D.P. Winchester, The eighth edition AJCC cancer staging manual: Continuing to build a bridge from a population-based to a more “personalized” approach to cancer staging, *CA Cancer J Clin* **67** (2017), 93–99.
- [19] O. Dworak, L. Keilholz and A. Hoffmann, Pathological features of rectal cancer after preoperative radiochemotherapy, *Int J Colorectal Dis* **12** (1997), 19–23.
- [20] S. Anfossi, A. Babayan, K. Pantel and G.A. Calin, Clinical utility of circulating non-coding RNAs – an update, *Nature Reviews Clinical Oncology* **15** (2018), 541.
- [21] P. Ranji, T.S. Kesejini, S. Saedikhoo and A.M. Alizadeh, Targeting cancer stem cell-specific markers and/or associated signaling pathways for overcoming cancer drug resistance, *Tumor Biology* **37** (2016), 13059–13075.
- [22] Ó. Rapado-González, A. Álvarez-Castro, R. López-López, J. Iglesias-Canle, M.M. Suárez-Cunqueiro and L. Muínelo-Romay, Circulating microRNAs as promising biomarkers in colorectal cancer, *Cancers* **11** (2019), 898.
- [23] Y. Shi, X. Zhang, X. Tang, P. Wang, H. Wang and Y. Wang, MiR-21 is continually elevated long-term in the brain after exposure to ionizing radiation, *Radiation Research* **177** (2011), 124–128.
- [24] J. Li, Y. Liu, C. Wang, T. Deng, H. Liang, Y. Wang, D. Huang, Q. Fan, X. Wang and T. Ning, Serum miRNA expression profile as a prognostic biomarker of stage II/III colorectal adenocarcinoma, *Scientific Reports* **5** (2015), 1–13.
- [25] X.X. Pu, G.L. Huang, H.Q. Guo, C.C. Guo, H. Li, S. Ye, S. Ling, L. Jiang, Y. Tian and T.Y. Lin, Circulating miR-221 directly amplified from plasma is a potential diagnostic and prognostic marker of colorectal cancer and is correlated with p53 expression, *Journal of Gastroenterology and Hepatology* **25** (2010), 1674–1680.
- [26] A. Sebio, L. Pare, D. Paez, J. Salazar, A. Gonzalez, N. Sala, E. del Río, M. Martín-Richard, M. Tobeña and A. Barnadas, The

- 721 LCS6 polymorphism in the binding site of let-7 microRNA 759
 722 to the KRAS 3'-untranslated region: Its role in the efficacy 760
 723 of anti-EGFR-based therapy in metastatic colorectal cancer 761
 724 patients, *Pharmacogenetics and Genomics* **23** (2013), 142– 762
 725 147. 763
- 726 [27] A. Pagliuca, C. Valvo, E.D. Fabrizio, S. Di Martino, M. Biffoni, 764
 727 D. Runci, S. Forte, R. De Maria and L. Ricci-Vitiani, Analysis 765
 728 of the combined action of miR-143 and miR-145 on onco- 766
 729 genic pathways in colorectal cancer cells reveals a coordinate 767
 730 program of gene repression, *Oncogene* **32** (2013), 4806–4813. 768
- 731 [28] J.V. Carter, N.J. Galbraith, D. Yang, J.F. Burton, S.P. Walker 769
 732 and S. Galandiuk, Blood-based microRNAs as biomarkers for 770
 733 the diagnosis of colorectal cancer: A systematic review and 771
 734 meta-analysis, *British Journal of Cancer* **116** (2017), 762–774. 772
- 735 [29] J.B. Weidhaas, I. Babar, S.M. Nallur, P. Trang, S. Roush, M. 773
 736 Boehm, E. Gillespie and F.J. Slack, MicroRNAs as poten- 774
 737 tial agents to alter resistance to cytotoxic anticancer therapy, 775
 738 *Cancer Research* **67** (2007), 11111–11116. 776
- 739 [30] Y. Sun, S. Shen, X. Liu, H. Tang, Z. Wang, Z. Yu, X. Li and M. 777
 740 Wu, MiR-429 inhibits cells growth and invasion and regulates 778
 741 EMT-related marker genes by targeting Onecut2 in colorectal 779
 742 carcinoma, *Molecular and Cellular Biochemistry* **390** (2014), 780
 743 19–30. 781
- 744 [31] K. Hur, Y. Toiyama, M. Takahashi, F. Balaguer, T. Nagasaka, 782
 745 J. Koike, H. Hemmi, M. Koi, C.R. Boland and A. Goel, 783
 746 MicroRNA-200c modulates epithelial-to-mesenchymal transi- 784
 747 tion (EMT) in human colorectal cancer metastasis, *Gut* **62** 785
 748 (2013), 1315–1326. 786
- 749 [32] M. Rokavec, H. Li, L. Jiang and H. Hermeking, The p53/ 787
 750 microRNA connection in gastrointestinal cancer, *Clinical and* 788
 751 *Experimental Gastroenterology* **7** (2014), 395. 789
- 752 [33] A.G. Renehan, L. Malcomson, R. Emsley, S. Gollins, A. Maw, 790
 753 A.S. Myint, P.S. Rooney, S. Susnerwala, A. Blower, M.P. 791
 754 Saunders, M.S. Wilson, N. Scott and S.T. O'Dwyer, Wait and- 792
 755 wait approach versus surgical resection after chemoradio- 793
 756 therapy for patients with rectal cancer (the OnCoRe study): 794
 757 A propensity-score matched cohort analysis, *Lancet Oncol* **17**
 758 (2016), 174–183.
- [34] C. Carames, I. Cristobal, V. Moreno, J.P. Marin, P. Gonzalez- 795
 Alonso, B. Torrejon, P. Minguez, A. Leon, J.I. Martin, R. Her- 796
 nandez, M. Pedregal, M.J. Martin, D. Cortes, D. Garcia-Olmo, 797
 M.J. Fernandez, F. Rojo and J. Garcia-Foncillas, MicroRNA- 798
 31 emerges as a predictive biomarker of pathological response 799
 and outcome in locally advanced rectal cancer, *Int J Mol Sci* 800
17 (2016). 801
- [35] G. Della Vittoria Scarpati, F. Falcetta, C. Carlomagno, P. 802
 Ubezio, S. Marchini, A. De Stefano, V.K. Singh, M. D'Incalci, 803
 S. De Placido and S. Pepe, A specific miRNA signature corre- 804
 lates with complete pathological response to neoadjuvant 805
 chemoradiotherapy in locally advanced rectal cancer, *Int J 806
 Radiat Oncol Biol Phys* **83** (2012), 1113–1119. 807
- [36] J. Yu, N. Li, X. Wang, H. Ren, W. Wang, S. Wang, Y. Song, 808
 Y. Liu, Y. Li, X. Zhou, A. Luo, Z. Liu and J. Jin, Circulating 809
 serum microRNA-345 correlates with unfavorable patho- 810
 logical response to preoperative chemoradiotherapy in locally 811
 advanced rectal cancer, *Oncotarget* **7** (2016), 64233–64243. 812
- [37] M. Campayo, A. Navarro, I. C. Benitez, S. Santasusagna, C. 813
 Ferrer, M. Monzo and I. C. Cera, miR-21, miR-99b and miR- 814
 375 combination as predictive response signature for preoper- 815
 ative chemoradiotherapy in rectal cancer, *PLoS One* **13** (2018), 816
 e0206542. 817
- [38] M. Marcuello, S. Duran-Sanchon, L. Moreno, J.J. Lozano, 818
 L. Bujanda, A. Castells and M. Gironella, Analysis of A 6- 819
 MiRNA signature in serum from colorectal cancer screening 820
 participants as non-invasive biomarkers for advanced adenoma 821
 and colorectal cancer detection, *Cancers* **11** (2019), 1542. 822
- [39] T. Machackova, V. Prochazka, Z. Kala and O. Slaby, Trans- 823
 lational potential of microRNAs for preoperative staging and 824
 prediction of chemoradiotherapy response in rectal cancer, 825
Cancers **11** (2019), 1545. 826
- [40] L. Imedio, I. Cristóbal, J. Rubio, A. Santos, F. Rojo and J. 827
 García-Foncillas, MicroRNAs in rectal cancer: Functional sig- 828
 nificance and promising therapeutic value, *Cancers* **12** (2020), 829
 2040. 830