

RESEARCH

Open Access



The effect of nutrition education for cancer prevention based on health belief model on nutrition knowledge, attitude, and practice of Iranian women

Bahareh Sasanfar^{1,2}, Fatemeh Toorang^{2,3}, Sahar Rostami², Maryam Zarif Yeganeh⁵, Maryam Lafzi Ghazi⁷, Monireh Sadat Seyyedsalehi^{2,8} and Kazem Zendehtdel^{2,4,6*}

Abstract

Background: In recent years, nutrition has received an increasingly important role in the etiology of cancer. Thus, public education about dietary factors associated with cancer risk or prevention could be an important intervention for cancer prevention, particularly in low- and middle-income countries where the burden of cancer is increasing rapidly and the access to care is limited. The age-standardized incidence of breast cancer was 35.8 among Iranian women in 2020. We aimed to study the effect of nutrition education on the knowledge, attitude, and practice of Iranian women towards dietary factors related to cancer.

Methods: In this interventional study, 229 women from public health centers were recruited and underwent three 75-min sessions of education based on the Health Belief Model (HBM). Participants were interviewed by trained interviewers using a validated and reproducible nutrition-related cancer prevention knowledge, attitude, and practice questionnaire (NUTCANKAP) questionnaire designed based on the HBM. Nutritional knowledge, attitude, and practice of participants were assessed through this questionnaire. Three 24-h dietary recalls (one weekend and two nonconsecutive weekdays) were also collected before and one month after the intervention.

Results: The mean age of the participants was 45.14 years, and the mean BMI was 27.2 kg/m². After the intervention, the participants had a higher intake of whole grain ($p = 0.03$) and a lower fat dairy ($p = 0.009$) and nuts ($p = 0.04$). However, the intake of high-fat dairy ($p = 0.001$) decreased after the intervention. We indicated significant differences in knowledge ($p < 0.001$) and nutritional practice scores ($p = 0.01$) after education. In addition, after the intervention, there were significant differences in the mean score of the HBM components, except for the perceived self-efficacy.

Conclusion: Participation in a nutrition education program positively impacted the knowledge and nutritional practices linked to cancer prevention.

Keywords: Health Belief model, Nutrition education, Nutrition knowledge, Cancer

Introduction

The burden of cancer continues to increase widely worldwide because of the population aging and increasing cancer-causing behaviors (e.g., unhealthy eating behavior, unhealthy food preparation) [1–3]. Based on the Global Cancer Observatory (GCO), which is the official cancer

*Correspondence: kzendeht@tums.ac.ir

² Cancer Research Center, Cancer Institute of Iran, Tehran University of Medical Sciences, P.O.Box: 13145158, Tehran, Iran
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

statistics of the International Agency for Research on Cancer (IARC), 29.5 million new cancer cases will be diagnosed worldwide in 2040 [2].

Primary prevention, including changing lifestyle and environmental interventions, has been illustrated as a key cancer control strategy for reducing this burden [4]. Previous research suggests that a combination of physical activity, having a healthy body weight, and a healthy diet could prevent one-third of cancers [5, 6]. Diet is a modifiable risk factor that can influence the risk of cancer. Several studies have investigated the relationship between dietary components, including fruit and vegetable, meat and processed meat, fiber intake, and the risk of cancer [7–9]. These studies illustrated that higher consumption of red and processed meat might increase breast cancer risk. However, adherence to a prudent dietary pattern containing high fruits, vegetables, and fibers might decrease breast cancer risk [10]. Furthermore, greater levels of nutrition knowledge have been linked to higher health literacy, better management of chronic diseases, and lower health costs [11]. Changing of attitude and practice has increasingly been used in nutrition education to improve intervention efficacy. Nutrition education programs could help to increase nutrition knowledge and improve dietary behaviors which may reduce the incidence of many chronic diseases including, cancer, diabetes, and cardiovascular disease [12]. Sullivan et al. illustrated that a nutrition education program strengthened nutrition-related cancer prevention attitudes among low-income African American women [13]. Another study assessed the effects of education on dietary behavior and showed that education plans based on HBM could change nutritional beliefs and behaviors for colorectal cancer prevention [14]. Regarding dietary intake behavior, an ecologic study and a meta-analysis of prospective studies found a positive relation between habitual salt intake and risk of gastric cancer [15, 16].

The Health Belief Model (HBM) is one of the most recommended models in the field of nutrition education programs [17]. This model describes the risks of unhealthy behaviour and the related and understands their susceptibility to adverse outcomes of their feelings and can be used as motivation to reduce risks [18]. This model includes five components: perceived susceptibility (i.e., the level in which a person knows his sensitiveness about a disease), perceived severity (i.e., the perceptions of the person about the severity of the disease), perceived benefits (i.e., the person's understanding about the advantages of the preventive behavior), perceived barriers (i.e., each healthy behavior and practice may encounter some barriers and problems), and performance guides (i.e., stimulations, which facilitate decision-making) [19–21]. Some studies have stated the benefits of using this model

in different health education programs [22, 23]. The use of the HBM is appropriate for myriad preventive health behaviors among both men and women; mainly, it has been a directive to researchers studying women's health issues [24]. In addition, nutrition researchers have frequently applied the HBM in cancer prevention education.

To our knowledge, there are limited studies about the effect of nutrition education on knowledge, attitude and practice for cancer prevention in Middle-Eastern countries. Therefore, we designed a prospective study in Iran to: (1) Assess the effect of nutrition education based on the HBM on the nutritional knowledge, attitude, and practice (KAP); (2) Recognize perceived barriers to the adherence of eating behaviors related to cancer prevention; (3) Change in nutritional behavior including food choice and methods of food processing that are associated with cancer prevention.

Subjects and methods

There were 229 participants who were visiting public health centers. We invited the women to participate in the study through flyers, posters, and introduction of this study on social media. We conducted an interventional study and used a one-group pretest–posttest design, through convenience sampling in 2017–2018, to evaluate the impact of nutrition education on the knowledge, attitude, and practice of women referring to the public health centers of Tehran University of Medical Science, located in Iran. The participating women were interviewed by trained interviewers using a validated and reproducible the nutrition-related cancer prevention knowledge, attitude and practice 36 questionnaire (NUTCANKAP) [25]. A 24-h dietary recall was conducted by phone on three different days, including two non-consecutive weekdays and a weekend before and one month after the intervention for all participants. The NUTCANKAP questionnaire was designed based on the HBM and consisted of three sections: A. knowledge (10 questions), B. attitude (27 questions; including 11 questions on perceived susceptibility, four questions on perceived severity, four questions on perceived benefits, four questions on perceived self-efficacy, and four questions on perceived barriers), and C. practice (16 questions). Correct answers in the knowledge section were given a score of 1. Incorrect answers, don't know-answers, and blanks were assigned a score of zero. The total raw scores of knowledge ranged from 0 to 10. The attitude section was evaluated by a Likert scale ranging from 1 as least desirable to 5 as most desirable, or vice versa. In the items related to the practice domain, correct food choices received a score of 1, and incorrect or blank responses were regarded as zero (Table 1). We calculated intakes of energy and all consumed foods through three recalls and then converted them to grams by a program

Table 1 Summary of item content and scoring of the NUTCANKAP questionnaire

Domain/constituent	No. of item	Example question	Scoring
Knowledge of cancer risk factors	10	Less intake of fried food can help to cancer prevention Higher intake of salt was associated with a risk of cancer	1 = Correct answer 0 = Incorrect or don't know-answer
Attitude towards cancer prevention	27	Method and time of food storage is important in cancer prevention	5 = Strong agree
Perceived susceptibility	11	Treatment cost are high for cancer	4 = Agree
Perceived severity	4	Sufficient vitamin D status may helps to cancer prevention	3 = Don't know
Perceived benefits	4	I can consumption moldy food after removing the mold from surface of food and heating it	2 = Disagree
Perceived self-efficacy	4	Flatulence from beans is a barrier to consuming them	1 = Strongly disagree (or vice versa)
Perceived barriers	4		
Practice for cancer prevention	16	I choose fruit juice rather than cola for beverage I choose boiling process rather than frying for food preparation	1 = Correct food choices 0 = Incorrect or blank responses

made by the authors in Microsoft Access. The nutrient composition of consumed foods was determined based on the USDA food composition database modified for Iranian foods. Participants who met the inclusion criteria were literate women aged 19–70 who had an available phone number for follow-up. Exclusion criteria included not being interested in continuing the study or being on dietary restrictions. Written informed consent was obtained from each participant.

After filling out the questionnaire, the educational program was performed in three 75-min sessions. Subjects were also given a book on cancer prevention through healthy nutrition. The educational program was designed based on the components of the HBM and pretest results was conducted through live lectures, collaborative question-answering methods, group discussions, and visual education materials such as slide shows. In the first session, the health educator informed them about cancer, potential risk factors, obesity and cancer, and healthy and unhealthy foods concerning cancer, through presentation slides. In the second session, the health educators held group discussions about the topics of previous session, types of cooking methods, and cooking dish. The educators also tried to promote attitude toward behavior in the participants. Information on food nutrition labels and the five food groups based on the food pyramid for reforming nutritional behavior was discussed in the third session [26]. In addition, a woman who had lost her first- or second-degree relative (s) due to cancer was invited to talk about the severity of the consequences of the disease. All procedures involving human subjects/patients were approved by the Research Ethics Committee of Tehran University of Medical Science (code: 28,614).

Finally, data were analyzed by STATA version 14 (State Corp., College Station, TX). The chi-square test and t-test were used for qualitative and continuous variables, respectively. Multivariable logistic regression was used to estimate the association between KAP scores and age,

educational and socioeconomic status. Age and socioeconomic were controlled as covariates.

Results

The mean age of the participants was 45.14 years (standard deviation = 10.16, range from 20 to 70), and the mean BMI was 27.2 kg/m². The majority of subjects were overweight (37.33%) and had a diploma (65.28%). Table 2 shows the association between KAP scores and age, educational, and socioeconomic status before the

Table 2 The association of studied KAP scores before the intervention between age, educational and socioeconomic groups

	Knowledge	Attitude	Practice
Age ¹	OR (CI)	OR (CI)	OR (CI)
20–41	Reference	Reference	Reference
42–50	0.56 (0.27–1.16)	1.42 (0.69–2.92)	1.07 (0.52–2.18)
51–69	0.86 (0.41–1.81)	1.46 (0.70–3.05)	1.84 (0.87–3.90)
†P for trend	0.48	0.33	0.13
Educational status ²			
Primary	Reference	Reference	Reference
Diploma	3.98 (1.21–13.10)	1.50 (0.52–4.30)	2.49 (0.85–7.29)
University	5.49 (1.47–20.39)	1.14 (0.33–3.35)	3.67 (1.09–12.37)
†P for trend	0.01	0.79	0.04
Socioeconomic status ²			
Low	Reference	Reference	Reference
Medium	1.84 (0.87–3.90)	1.33 (0.63–2.81)	1.22 (0.58–2.56)
High	1.07 (0.51–2.21)	0.80 (0.39–1.66)	1.11 (0.53–2.31)
†P for trend	0.76	0.52	0.73

Significant P value (< 0.05) was bolded

¹ adjusted for Socioeconomic status

² adjusted for age

†Obtained from logistic regression. OR has been computed by considering subjects over medium scores as one and lower medium scores as zero

intervention. Women with university education had higher knowledge ($p=0.01$) and nutritional practice ($p=0.04$) scores than those with primary education. However, no significant differences were observed across age and socioeconomic groups. After the intervention, women reduced intake of carbohydrate ($p=0.008$), total protein ($p=0.03$), animal protein ($p=0.05$), vegetable fat ($p=0.01$), saturated fatty acid ($p=0.0002$), monounsaturated fatty acid ($p=0.04$), Cobalamin ($p=0.01$), Iron ($p=0.01$), and Selenium ($p=0.006$) (Table 3). In addition, after the intervention subjects had higher intake of whole grain ($p=0.03$), low fat dairy ($p=0.009$), and nuts ($p=0.04$). However, the intake of high-fat dairy ($p=0.001$) decreased (Table 4) after education. Comparisons of the scores of knowledge, attitude (perceived susceptibility, perceived severity, perceived benefits, perceived barriers, and perceived self-efficacy) and nutritional practice, before and after the education are presented in Table 5. We indicated significant differences in knowledge ($p<0.001$) and nutritional practice scores ($p<0.001$) following education. Moreover, after

the intervention, there were significant differences in attitude ($p<0.001$). Therefore, the mean score of knowledge, attitude and nutritional practice significantly increased after the intervention. We found a significant association between the improvements of attitude score after the intervention ($P_{\text{trend}}=0.04$) and the level of education (Table 6). The association was also significant for the specific question of knowledge about BMI and the level of education (OR=6.27; 95% CI=1.72–22.7, $P_{\text{trend}}=0.001$) and socioeconomic status (OR=0.39; 95% CI=0.18–0.83, $P_{\text{trend}}=0.01$). In addition, we found a significant association between sing food labels and the level of education (OR=6.07; 95% CI=1.71–21.5, $P_{\text{trend}}=0.006$).

Comparison of the scores obtained from question 17 of nutritional practice before and after the intervention is shown as a sample in Table 7.

Discussion

This study is the first study that used the HBM in nutrition education for cancer prevention in Iran. We concluded that the application of the HBM in nutrition

Table 3 Comparison of macro- and micronutrient intake before and after the intervention

	Before intervention	After intervention	<i>p</i> -value
Energy (kcal)	2200.5 ± 62.0	2117.7 ± 60.0	0.11
Carbohydrate (g/d)	295.29 ± 52.59	282.72 ± 46.54	0.008
Total protein (g/d)	66.37 ± 15.52	62.80 ± 21.87	0.03
Animal protein (g/d)	35.27 ± 24.87	31.35 ± 18.13	0.05
Vegetable protein (g/d)	37.98 ± 8.60	37.85 ± 8.80	0.44
Total fat (g/d)	60.22 ± 21.04	59.22 ± 21.14	0.32
Animal fat (g/d)	27.09 ± 14.03	25.33 ± 14.99	0.13
Vegetable fat (g/d)	40.09 ± 21.97	35.37 ± 19.80	0.01
Saturated fatty acid (g/d)	20.38 ± 9.06	17.15 ± 10.34	0.0002
Monounsaturated fatty acid (g/d)	19.95 ± 8.03	18.56 ± 7.63	0.04
Polyunsaturated fatty acid (g/d)	16.62 ± 12.27	16.61 ± 11.10	0.49
Cholesterol (mg)	160.00 ± 116.38	156.07 ± 86.83	0.36
Fiber (g/d)	20.26 ± 10.88	19.90 ± 7.17	0.32
Vitamin C	159.84 ± 131.67	154.82 ± 100.66	0.32
Vitamin A (µg)	855.33 ± 792.55	839.79 ± 589.52	0.40
Thiamin (mg)	1.60 ± 0.35	1.55 ± 0.36	0.10
Riboflavin (mg)	1.35 ± 0.49	1.30 ± 0.48	0.09
Niacin (mg)	17.62 ± 27.32	15.84 ± 33.83	0.30
Pyridoxin (mg)	1.43 ± 0.49	1.36 ± 0.46	0.06
Cobalamin (µg)	1.22 ± 2.59	0.21 ± 3.66	0.001
Folate (µg)	279.78 ± 10.63	282.61 ± 11.15	0.58
Iron (mg)	20.42 ± 9.38	18.78 ± 7.69	0.01
Calcium (mg)	899.70 ± 342.19	865.14 ± 347.28	0.09
Zinc (mg)	7.59 ± 3.02	7.27 ± 1.73	0.11
Selenium (µg)	80.48 ± 3.10	77.67 ± 3.02	0.006

Significant *P* value (< 0.05) was bolded

Values are mean (SD)

Table 4 Comparison of food group intake before and after the intervention

Food group	Before intervention	After intervention	p-value
Wholegrain	24.92 ± 64.49	37.17 ± 59.27	0.03
Refined grain	217.57 ± 114.63	231.02 ± 131.52	0.14
Low-fat dairy	136.27 ± 152.25	166.68 ± 146.31	0.009
Medium fat dairy	27.78 ± 80.76	30.32 ± 71.53	0.37
High-fat dairy	135.37 ± 191.41	83.37 ± 150.66	0.001
Meat	70.117 ± 108.27	65.78 ± 73.18	0.33
Processed meat	10.77 ± 39.30	7.91 ± 31.82	0.22
Fish	8.05 ± 21.57	10.49 ± 29.40	0.16
Vegetables	318.32 ± 189.65	313.12 ± 229.15	0.40
Fruits	522.79 ± 297.39	560.79 ± 275.61	0.07
Fruit juice	34.27 ± 86.58	30.90 ± 67.18	0.33
Beans	36.06 ± 50.87	32.40 ± 54.27	0.27
Sweet and desserts	15.65 ± 54.38	16.67 ± 42.22	0.42
Hydrogenated fat	4.14 ± 17.39	4.33 ± 9.54	0.45
Animal fat	3.78 ± 8.17	4.67 ± 11.70	0.18
Olive and other vegetable oil	5.20 ± 17.72	5.31 ± 12.37	0.47
Egg	9.82 ± 27.61	9.09 ± 27.08	0.40
Nuts	4.27 ± 23.44	8.95 ± 28.71	0.04

Significant P value (< 0.05) was bolded

Values are mean (SD)

Table 5 Comparison of studied KAP scores before and after the intervention

Variable	Before intervention	After intervention		p-value
	Mean (± SD)	Mean (± SD)	Differences(± SD)	
Knowledge†	33.83 (10.05)	41.12 (7.32)	8.14 (9.29)	< 0.001
Attitude†	62.22 (18.45)	71.82 (19.23)	9.59 (20.9)	< 0.001
Perceived susceptibility	44.71 (5.13)	45.96 (4.60)	1.24 (3.45)	0.001
Perceived severity	65.54 (21.62)	77.12 (16.94)	11.57 (23.13)	< 0.001
Perceived benefits	59.61 (16.72)	65.23 (13.52)	5.62 (19.33)	0.0002
Perceived barriers	77.27 (40.05)	89.93 (29.91)	12.66 (42.99)	0.0002
Perceived self-efficacy	78.95 (17.23)	84.40 (18.61)	5.44 (21.74)	0.001
Practice†	69.56 (15.92)	79.19 (12.83)	9.62 (15.14)	< 0.001

SD standard deviation

P-values were determined by the T-test

†The values are shown in the scale of 0–100

education for cancer prevention could result in promoting the level of knowledge, attitude, and nutritional practice among Iranian women. Our results showed a low level of knowledge about cancer causes, protective nutrients and those lowering the risk of cancer, healthy cooking methods, food guide pyramid, and healthy cooking dish before the intervention. The score of knowledge was higher among women with a university education than other groups before the intervention ($p < 0.01$). Knowledge of participants about normal BMI range doubled

after education (0.14 vs. 0.28). However, the mean score of knowledge significantly increased after the intervention. There was a non-significant increasing trend in the difference in knowledge score between age and education. However, the before-after differences in knowledge and practice among participants with a high socioeconomic status were lower than those at the higher socioeconomic level. This may suggest that higher economic levels do not necessarily reflect greater awareness. Multiple levels of influence affect an individual's food choice.

Table 6 The association of differences of KAP scores between age, educational and socioeconomic groupst

	Differences in knowledge score		Differences in attitude score	Differences in practice score		
	Overall	Q3 [†]	Overall	Overall	Q14 [†]	Q17 ^{††}
Age ¹	OR (CI)	OR (CI)	OR (CI)	OR (CI)	OR (CI)	OR (CI)
20–41	Reference	Reference	Reference	Reference	Reference	Reference
42–50	0.87 (0.42–1.80)	0.97 (0.46–2.01)	0.97 (0.46–2.04)	1.13 (0.55–2.31)	0.79 (0.38–1.63)	0.96 (0.41–2.23)
51–69	1.29 (0.60–2.77)	1.59 (0.74–3.39)	0.94 (0.44–1.99)	0.84 (0.41–1.74)	1.40 (0.65–3.03)	1.06 (0.42–2.65)
[†] P for trend	0.54	0.25	0.87	0.69	0.46	0.91
Educational status ²						
Primary	Reference	Reference	Reference	Reference	Reference	Reference
Diploma	1.27 (0.44–3.60)	1.21 (0.42–3.46)	1.09 (0.38–3.10)	1.23 (0.43–3.48)	3.15 (1.06–9.29)	0.76 (0.23–2.48)
University	2.17 (0.66–7.11)	6.27 (1.72–22.7)	2.75 (0.80–9.46)	1.72 (0.53–5.55)	6.07 (1.71–21.57)	0.52 (0.12–2.09)
[†] P for trend	0.13	0.001	0.04	0.29	0.006	0.36
Socioeconomic status ²						
Low	Reference	Reference	Reference	Reference	Reference	Reference
Medium	0.89 (0.41–1.89)	0.71 (0.33–1.52)	0.96 (0.45–2.06)	1.17 (0.56–2.44)	0.69 (0.32–1.49)	1.19 (0.48–2.90)
High	0.60 (0.28–1.28)	0.39 (0.18–0.83)	0.99 (0.46–2.11)	0.92 (0.44–1.90)	0.57 (0.26–1.22)	1.84 (0.77–4.42)
[†] P for trend	0.21	0.01	0.98	0.82	0.15	0.16

Significant P value (< 0.05) was bolded

¹ adjusted for socioeconomic

² adjusted for age

[†] Q3 = What is the normal range of BMI for adults?

^{††} Q14 = Do you use food labels when choosing food?

^{†††} Q17 = Do you use high-fat dairy?

[†] Obtained from logistic regression. OR has been computed by considering subjects over medium scores as one and lower medium scores as zero

Table 7 Comparison of question 17 scores of practice before and after the intervention (question 17: which of the following do you do?)

	Before	After	P*
The overall score of practice question 17	0.24 ± 0.21	0.42 ± 0.36	< 0.001
I put my hands and feet for a few minutes against the sun to make vitamin D	0.26 ± 0.44	0.37 ± 0.48	0.003
I check my iron and folic acid levels under the guidance of a nutrition consultant	0.36 ± 0.48	0.64 ± 0.47	0.0000
I will reduce the consumption of simple sugar like sweets	0.74 ± 0.43	0.92 ± 0.26	0.0000
I don't use soda even zero types	0.53 ± 0.50	0.70 ± 0.45	0.0002
I check my weight every week	0.41 ± 0.49	0.53 ± 0.50	0.001
I use low-fat dairy instead of high-fat dairy	0.70 ± 0.45	0.77 ± 0.41	0.04

* Obtained from independent student's t-test

Biological and cultural influences such as taste, sex, and age may have significant effects on food consumption [27, 28]. This suggests that food consumption is not necessarily associated with the cost [29]. on the other hand, families have to pay rent, clothes, and transport in addition to buying food. Therefore, only a tiny part of their income is allocated to food.

Ahn et al. [30] showed that nutritional education positively impacted dietary habits and nutritional knowledge in older adults. A group of researchers in the USA conducted a program to teach children about cancer and

cancer control behaviors and found it successful in promoting the knowledge about cancer risk factors, forming a positive attitude towards cancer risk factors, and increasing cancer control behaviors among students [31]. It should be kept in mind that nutrition literacy enables people to use written information related to health. Therefore, increasing nutrition knowledge has a protective effect against diseases. A systematic review illustrated that the majority of studies reported a significant association between nutrition knowledge and dietary intake [11].

We used the HBM to increase the impact of nutrition education. The HBM would seem to be used widely for communication research [32], and has been suggested [33] and approved [34] as a model for nutrition education. In our study, the score of the HBM constructs, including perceived susceptibility, perceived severity, perceived benefits, perceived barriers, and perceived self-efficacy increased following a nutrition education program. Additionally, in this study, the score of perceived barriers increased after the intervention. This implies that women recognized the barriers and would try to resolve them. Similarly, an interventional study on gastric cancer among 84 Iranian housewives showed that the intervention group based on the HBM model showed significantly higher scores after the education [35]. Attitude scores showed a decreasing trend between age, educational, and socioeconomic status before the intervention. However, the difference in attitude scores had an increasing trend after the intervention unless the socioeconomic status. Meanwhile, an interventional study on 157 African American women on nutrition-related cancer prevention showed that attitudes improved after a nutrition education program [13].

This study illustrated that nutrition education program based on the HBM has a positive effect on food choices among women. We assessed this change through a questionnaire and three recalls and found that the score of nutritional practice increased after the intervention.

The question score on food labeling usage increased by 82%, and the question score of high-fat dairy usage increased by 33% after the education. Therefore, participants had a better food choice following nutrition education. An increase in whole grain, low-fat dairy, and nuts was also found after the intervention. These food groups compose of nutrients protecting against cancer [36]. In addition, we found a decrease in carbohydrate, total protein, animal protein, vegetable fat, saturated fatty acid, monounsaturated fatty acid, Cobalamin, Iron, Selenium, and high-fat dairy. Studies illustrated that carbohydrate intake is positively associated with cancer via insulin and the related hormone, IGF-1 [37–39]. We found significant associations for the specific question of knowledge on BMI and level of education (OR = 6.44) and socioeconomic status (OR = 0.39). Similarly, differences in knowledge, attitude and nutritional practice scores between socioeconomic groups showed a decreasing trend after the intervention. Jing Wu et al. [40] suggested that a higher intake of total red meat, fresh red meat, processed meat, and high-fat dairy may be risk factors for breast cancer. Regarding fatty acids, various studies have demonstrated that polyunsaturated fatty acids possess a therapeutic role against certain types of cancer [41]. In contrast, the intake of saturated fatty acids

has been linked to cancer [42]. A study evaluated medical students' knowledge about the association between dietary factors and the risk of cancer and indicated that diet-disease knowledge was higher among those who had a higher dietary fiber intake [43]. In our study, increasing in knowledge score was seen toward vegetables and fruits consumption, but it was not achieved in nutritional practice scores. The barriers to low fruit intake in participants were determined through the questionnaire. Limited budget was mentioned by 13.4% of participants as the main barrier to fruit consumption. Around 10% of the participating women believed that preparing fruits is time-consuming which could be a barrier to fruit intake, and 2.59% limited their fruit consumption due to digestive problems. A small part of our study sample (1.04%) cut fruit intake because they believe fruits are contaminated with toxins. In addition, participants reported a lack of vegetable consumption due to difficulty in preparation (18.75%), cost (2.07%), lack of irrigation with safe water (8.29%), or digestive problems (8.29%).

A study conducted in Northwest of Iran illustrated that the food habits of East-Azerbaijan people in the last two decades increase the risk of gastric cancer and suggested performing nutrition education for a healthy diet [44]. In the Golestan cohort study, the incidence of esophageal cancer was associated with nutrient intake and dietary behaviors such as polycyclic aromatic hydrocarbons and drinking hot tea [45]. Other Iranian studies indicated that nutrition-related attitudes were positively correlated with the dietary practices of breast cancer prevention [46]. Since Iranian unique dietary habits are modifiable by education and with regards to the burden of high health system costs of cancer imposed on patients and the government, the application of education programs would be cost-effective.

This study was limited to the intervention group. Our study was done before and one month after the intervention, which only showed the short-term effects of the intervention. The study population was limited to females. Women have a critical role in food choices and nutrition education of children in the family. Due to strong linkage between maternal education and children's health, we conducted this study among women. However, the results of this study cannot be generalized to men, and additional research among Iranian men is needed. Some confounding variables such as personality characteristics, mental health, and media might have affected the outcome, which was not assessed. According to the differences in scores between different age groups, a different educational approach may have to be applied to each age group. The strengths of the current study include large sample size, recruiting participants from various areas of Tehran, and the use of visual education

materials. We also used a validated instrument to measure educational intervention and assessed food practice by collecting dietary recalls, which have not been done by many studies [25].

In conclusion, this study showed that a nutrition education program based on the HBM had a positive impact on the knowledge and nutritional practice of Iranian women. Considering the cost-effectiveness of educational programs compared to treatment services, applying health education programs can highly promote public health.

Acknowledgements

We would like to express our special thanks to the participants and healthcare executives in public health centers, without whom the study would not have been possible.

Author contributions

BS, FT, and MZ participated in designing the study, analysis, and drafting the initial version. FT helped in data analysis. BS implemented comments and suggestions from the co-authors. SR, MG, and MS helped in applying of the study. KZ contributed to the conception, design, and data analysis. All authors reviewed the final version of the manuscript. KZ supervised the study.

Funding

This study was financially supported by Cancer Research Center of Tehran University of Medical Science (no. 94-01-51-28614).

Availability data and materials

Data is available on reasonable request by corresponding author (Dr. Kazem Zندهدل).

Declarations

Ethical approval and consent to participate

This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by Tehran University of Medical Science (code: 28614). Written informed consent was obtained from all subjects/patients.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. ²Cancer Research Center, Cancer Institute of Iran, Tehran University of Medical Sciences, P.O.Box: 13145158, Tehran, Iran. ³Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran. ⁴Cancer Biology Research Center, Cancer Institute of Iran, Tehran University of Medical Sciences, Tehran, Iran. ⁵Department of Community Nutrition Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ⁶Breast Diseases Research Center, Cancer Institute of Iran, Tehran University of Medical Sciences, Tehran, Iran. ⁷Department of Exercise Physiology, Central Tehran Branch, Islamic Azad University, Tehran, Iran. ⁸Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy.

Received: 2 November 2021 Accepted: 27 May 2022

Published online: 07 June 2022

References

1. Mayne ST, Playdon MC, Rock CL. Diet, nutrition, and cancer: past, present and future. *Nat Rev Clin Oncol*. 2016;13(8):504.
2. Golzar A, et al. Food preparation methods, drinking water source and esophageal squamous cell carcinoma in the high risk area of golestan, Northeast Iran. *European J Cancer Prev Official J European Cancer Prev Organ (ECP)*. 2016;25(2):123.
3. Patel A, et al. Role of nutritional factors in pathogenesis of cancer. *Food Quality Safe*. 2018;2(1):27–36.
4. Zhang Y-B, et al. Combined lifestyle factors, incident cancer, and cancer mortality: a systematic review and meta-analysis of prospective cohort studies. *Br J Cancer*. 2020;122(7):1085–93.
5. Kohler LN, et al. Adherence to diet and physical activity cancer prevention guidelines and cancer outcomes: a systematic review. *Cancer Epidemiol Prev Biomarkers*. 2016;25(7):1018–28.
6. Rock CL et al. American cancer society guideline for diet and physical activity for cancer prevention. *CA: Cancer J Clin*. 2020;70(4):245–271.
7. Garcia-Larsen V, et al. Dietary patterns derived from principal component analysis (PCA) and risk of colorectal cancer: a systematic review and meta-analysis. *European J Clin Nutr*. 2018. <https://doi.org/10.1038/s41430-018-0234-7>.
8. Farvid MS, et al. Consumption of red and processed meat and breast cancer incidence: A systematic review and meta-analysis of prospective studies. *Int J Cancer*. 2018;143(11):2787–99.
9. McRae MP. The benefits of dietary fiber intake on reducing the risk of cancer: an umbrella review of meta-analyses. *J Chiropr Med*. 2018;17(2):90–6.
10. Sasanfar B, et al. Adherence to plant-based dietary pattern and risk of breast cancer among Iranian women. *Eur J Clin Nutr*. 2021;75(11):1578–87.
11. Spronk I, et al. Relationship between nutrition knowledge and dietary intake. *Br J Nutr*. 2014;111(10):1713–26.
12. Hutchinson A, Wilson B, Wilson C. Cancer prevention through nutrition education in the workplace a review of the literature. *Int J Cancer Prev*. 2010;3(3):111–36.
13. Sullivan HW, Klassen AC. Nutrition-related cancer prevention attitudes in low-income women. *Prev Med*. 2007;45(2–3):139–45.
14. Hatami T, et al. Effect of multimedia education on nutritional behaviour for colorectal cancer prevention: an application of health belief model. *Malaysian J Med Sci MJMS*. 2018;25(6):110.
15. D'Elia L, et al. Habitual salt intake and risk of gastric cancer: a meta-analysis of prospective studies. *Clin Nutr*. 2012;31(4):489–98.
16. Park B, et al. Ecological study for refrigerator use, salt, vegetable, and fruit intakes, and gastric cancer. *Cancer Causes Control*. 2011;22(11):1497–502.
17. Khoramabadi M, et al. Effects of education based on health belief model on dietary behaviors of Iranian pregnant women. *Global J Health Sci*. 2016;8(2):230.
18. Gilmore GD. Needs and capacity assessment strategies for health education and health promotion-BOOK ALONE. Jones & Bartlett Publishers; 2011.
19. Martin LR, Haskard-Zolnieriek KB, DiMatteo MR. Health behavior change and treatment adherence: evidence-based guidelines for improving healthcare. USA: Oxford University Press; 2010.
20. Rezapour B, Mostafavi F, Khalkhali H. "Theory Based Health Education: Application of Health Belief Model for Iranian Obese and Overweight Students about Physical Activity" in Urmia Iran. *Int J Prev Med*. 2016. <https://doi.org/10.4103/2008-7802.191879>.
21. Solhi M, et al. The application of the health belief model in oral health education. *Iran J Public Health*. 2010;39(4):114.
22. Ghaderi N, et al. Effect of education based on the Health Belief Model (HBM) on anemia preventive behaviors among iranian girl students. *Int J Pediatr*. 2017;5(6):5043–52.
23. Tavakoli HR, et al. Efficacy of HBM-based dietary education intervention on knowledge, attitude, and behavior in medical students. *Iranian Red Crescent Med J*. 2016. <https://doi.org/10.5812/ircmj.23584>.
24. Tanner-Smith EE, Brown TN. Evaluating the health belief model: a critical review of studies predicting mammographic and pap screening. *Soc Theory Health*. 2010;8(1):95–125.
25. Sasanfar B, et al. Development and validation of a knowledge, attitude, and practice questionnaire on nutrition-related cancer prevention for

- Iranian women. *J Res Med Sci Official J Isfahan Univ Med Sci*. 2019. https://doi.org/10.4103/jrms.JRMS_777_18.
26. Hart S, et al. Development of the "Recovery from Eating Disorders for Life" Food Guide (REAL Food Guide)-a food pyramid for adults with an eating disorder. *J Eat Disord*. 2018;6(1):1–11.
 27. Nettore IC, et al. Influences of age, sex and smoking habit on flavor recognition in healthy population. *Int J Environ Res Public Health*. 2020;17(3):959.
 28. Cattaneo C, et al. Cross-cultural differences in lingual tactile acuity, taste sensitivity phenotypical markers, and preferred oral processing behaviors. *Food Qual Prefer*. 2020;80: 103803.
 29. Middaugh AL, et al. Few associations between income and fruit and vegetable consumption. *J Nutr Educ Behav*. 2012;44(3):196–203.
 30. Ahn JA, Park J, Kim CJ. Effects of an individualised nutritional education and support programme on dietary habits, nutritional knowledge and nutritional status of older adults living alone. *J Clin Nurs*. 2018;27(9–10):2142–51.
 31. Ayers K, et al. Evaluation of the St. Jude cancer education for children program on cancer risk awareness, attitudes, and behavioral intentions among fourth-grade science students: comparisons between racially identifiable/high-poverty schools and racially diverse/affluent schools. *J Cancer Educ*. 2019. <https://doi.org/10.1007/s13187-019-1476-3>.
 32. Jones CL, et al. The health belief model as an explanatory framework in communication research: exploring parallel, serial, and moderated mediation. *Health Commun*. 2015;30(6):566–76.
 33. Vahedian-Shahroodi M, et al. The impact of health education on nutritional behaviors in female students: an application of health belief model. *Int J Health Promot Educ*. 2021;59(2):70–82.
 34. Cox DN, et al. Take Five, a nutrition education intervention to increase fruit and vegetable intakes: impact on consumer choice and nutrient intakes. *Br J Nutr*. 1998;80(2):123–31.
 35. Alidosti M, et al. An investigation on the effect of gastric cancer education based on health belief model on knowledge, attitude and nutritional practice of housewives. *Iran J Nurs Midwifery Res*. 2012;17(4):256.
 36. Aune D, et al. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *BMJ*. 2016. <https://doi.org/10.1136/bmj.i2716>.
 37. Fine EJ, Feinman RD. Insulin, carbohydrate restriction, metabolic syndrome and cancer. *Expert Rev Endocrinol Metab*. 2015;10(1):15–24.
 38. Amadou A, et al. Dietary carbohydrate, glycemic index, glycemic load, and breast cancer risk among Mexican women. *Epidemiology*. 2015;26(6):917–24.
 39. Sasanfar B, et al. Adherence to the low carbohydrate diet and the risk of breast cancer in Iran. *Nutr J*. 2019;18(1):86.
 40. Wu J, et al. Dietary protein sources and incidence of breast cancer: a dose-response meta-analysis of prospective studies. *Nutrients*. 2016;8(11):730.
 41. Nabavi SF, et al. Omega-3 polyunsaturated fatty acids and cancer: lessons learned from clinical trials. *Cancer Metastasis Rev*. 2015;34(3):359–80.
 42. Xia H, et al. Meta-analysis of saturated fatty acid intake and breast cancer risk. *Medicine*. 2015. <https://doi.org/10.1097/MD.0000000000002391>.
 43. Kendel Jovanović G, et al. Cancer and cardiovascular diseases nutrition knowledge and dietary intake of medical students. *Coll Antropol*. 2011;35(3):765–74.
 44. Somi MH, et al. Is there any relationship between food habits in the last two decades and gastric cancer in North-Western Iran. *Asian Pac J Cancer Prev*. 2015;16(1):283–90.
 45. Islami F, et al. Tea drinking habits and oesophageal cancer in a high risk area in northern Iran: population based case-control study. *BMJ*. 2009;338: b929.
 46. Raji Lahiji M, et al. Nutrition knowledge, attitudes, and practice towards breast cancer prevention among the female population of Iran University of medical science students. *Nutr Cancer*. 2019. <https://doi.org/10.1080/01635581.2019.1607410>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

